arXiv:quant-ph/9810069v1 23 Oct 1998

A RIGOROUS PATH-INTEGRAL FORMULA FOR QUANTUM SPIN DYNAMICS VIA PLANAR BROWNIAN MOTION

BERNHARD BODMANN

Department of Mathematics, University of Florida, 358 Little Hall, Gainesville, Fl 32611, USA

HAJO LESCHKE and SIMONE WARZEL

Institut für Theoretische Physik, Universität Erlangen-Nürnberg, Staudtstr. 7, D-91058 Erlangen, Germany

Adapting ideas of Daubechies and Klauder we derive a continuum path-integral formula for the time evolution generated by a spin Hamiltonian. For this purpose we identify the finite-dimensional spin Hilbert space with the ground-state eigenspace of a suitable Schödinger operator on $L^2(\mathbb{R}^2)$, the Hilbert space of square-integrable functions on the Euclidean plane $\mathbb{R}^2$, and employ the Feynman-Kac-Itô formula.

1 Introduction

Even 50 years after the appearance of Feynman's celebrated paper[1] which introduced the path-integral formalism into quantum theory in a heuristic but convincing manner, there is no general consensus on how to treat a quantum spin within this framework. To our knowledge, among the various approaches the only rigorous expression for the dynamics of a quantum spin in terms of an integral over continuous paths is due to Daubechies and Klauder.[2] They were able to write the coherent representation of the unitary time-evolution operator of a spin with definite quantum number as the ultra-diffusive limit of a well-defined integral over Brownian-motion paths on the two-sphere.

The present contribution summarizes a recent work[3], where we have shown that one may equally well employ Brownian-motion paths on the Euclidean plane. In this way a closer contact to symbolic continuum path-integral formulas widely discussed in the recent literature[4−7] is established. One may hope that the wealth of analytical tools associated with planar Brownian motion helps clarifying some subtle points there. The presented approach might also be of interest in the context of quantizing constrained systems.

2 Basic definitions, result, and comments

We consider a single **spin** with fixed **quantum number** $j \in \{0, 1/2, 1, 3/2, \ldots\}$. It is given by three operators $\mathcal{J}_+, \mathcal{J}_-$, and $\mathcal{J}_3$ which are viewed as acting on the $(2j+1)$-dimensional complex Hilbert space $\mathbb{C}^{2j+1}$ and obey the usual commutation relations $[\mathcal{J}_+, \mathcal{J}_-] = 2\mathcal{J}_3$, $[\mathcal{J}_3, \mathcal{J}_\pm] = \pm\mathcal{J}_\pm$.

Contribution to the conference proceedings:
Path integrals from peV to TeV - Firenze '98,
eds. V.Tognetti, R.Vaia, R.Casalbuoni, R.Giachetti, P.Verrucchi,
World Scientific, Singapore 1999

Non-normalized **coherent vectors**

$$|z\rangle := [(2j+1)/\pi]^{1/2} \left(1+|z|^2\right)^{-j-1} e^{z\mathcal{J}_+}|j,-j\rangle\,, \quad z \in \mathbb{C}\,, \tag{1}$$

in $\mathbb{C}^{2j+1}$ are parametrized by complex numbers. Henceforth, z^* will refer to their complex conjugates, $z_1 := (z+z^*)/2$ and $z_2 := (z-z^*)/2i$ to their real and imaginary parts. A normalized **spin-down vector** $|j,-j\rangle \in \mathbb{C}^{2j+1}$, obeying $\mathcal{J}_-|j,-j\rangle = 0$ and $\langle j,-j|j,-j\rangle = 1$, serves as the reference vector.

Every **spin Hamiltonian** $\mathcal{H}$, a (self-adjoint) operator on $\mathbb{C}^{2j+1}$, is polynomial in $\mathcal{J}_+, \mathcal{J}_-$, and $\mathcal{J}_3$ and may be written in **pseudo-diagonal form**

$$\mathcal{H} = \int_{\mathbb{C}} d^2z\, h(z)\, |z\rangle\langle z|\ . \tag{2}$$

Here the **contravariant symbol** h of $\mathcal{H}$ is a (real-valued) function on $\mathbb{C}$ which may be chosen bounded and continuous, and $d^2z := dz_1dz_2$ is the Lebesgue measure on the Euclidean plane $\mathbb{R}^2 := \mathbb{R}\times\mathbb{R} \cong \mathbb{C}$. In particular, the unit operator $\mathbb{1}$ on $\mathbb{C}^{2j+1}$ has the constant 1 as a contravariant symbol.

Furthermore, for given $z, z' \in \mathbb{C}$, $t > 0$, and $\nu > 0$ let $\mathbb{E}(\cdot)$ denote the probabilistic expectation with respect to the **two-dimensional Brownian bridge** with diffusion constant ν starting in $z = b(0)$ and arriving at $z' = b(t)$ a time t later.[8,9] As a Gaussian stochastic process with continuous paths $\{s \mapsto b(s) = b_1(s) + ib_2(s)\}_{s\in[0,t]}$ on $\mathbb{R}^2$ it is characterized by its means $\mathbb{E}(b_k(s)) = z_k + (z'_k - z_k)s/t$ and covariance functions $\mathbb{E}(b_k(r)b_l(s)) - \mathbb{E}(b_k(r))\mathbb{E}(b_l(s)) = 2\nu\delta_{kl}\,(\min\{r,s\} - rs/t)$; $k,l \in \{1,2\}$; $r,s \in [0,t]$. With the well-defined Wiener type of path integration $\int d\mu^{(\nu)}_{z,0;z',t}(b)\ (\cdot) := \exp\{-|z-z'|^2/4t\nu\}\,\mathbb{E}(\cdot)/4\pi t\nu$ the **coherent representation** $\langle z|\exp\{-it\mathcal{H}\}|z'\rangle$ of the (unitary) **spin time-evolution operator** $\exp\{-it\mathcal{H}\}$ may, for all $z, z' \in \mathbb{C}$ and $t > 0$, be expressed as an **ultra-diffusive limit** in the following sense

$$\boxed{\begin{aligned}\langle z|e^{-it\mathcal{H}}|z'\rangle = \lim_{\nu\to\infty} \int d\mu^{(\nu)}_{z,0;z',t}(b)\ \exp\left\{4(j+1)\nu\int_0^t \frac{ds}{(1+|b(s)|^2)^2}\right\}\\ \times\exp\left\{(j+1)\int_0^t ds\frac{\dot{b}(s)b^*(s) - \dot{b}^*(s)b(s)}{1+|b(s)|^2} - i\int_0^t ds\, h(b(s))\right\}\end{aligned}}\ . \tag{3}$$

This is the main result of Ref. 3. Here the second integral in the exponent is a purely imaginary stochastic (line) integral, which can be understood in the sense of Fisk and Stratonovich and to which one is therefore allowed to apply the rules of ordinary calculus, although the time derivative $\dot{b}$ does not exist.[9]

Several comments apply:

- Keeping $t > 0$ but replacing h by $-h$ or $-ih$ in (3) yields analogous path-integral expressions for the coherent representation of the inverse spin time-evolution $\exp\{it\mathcal{H}\}$ or the **spin Boltzmann operator** $\exp\{-t\mathcal{H}\}$.

- The flat-space Wiener-regularized path-integral expression (3) for the spin time-evolution operator is an alternative to a result given in Ref. 2 relying on spherical Brownian motion. Contrary to what one might expect, equation (3) cannot be obtained from this result merely by stereographically projecting the paths from the sphere onto the (extended) Euclidean plane. In particular, unlike in Ref. 2, the regularization used in (3) breaks part of the $SU(2)$ symmetry when realized on $\mathbb{C}$. The full symmetry is restored only in the limit.

- In order to make contact with the Wiener-regularized path-integral expression associated with the dynamics of a canonical degree of freedom, also proved in Ref. 2, one has to contract the $su(2)$ algebra to the Heisenberg-Weyl algebra by taking the **high-spin limit** $j \to \infty$.[10] More precisely, in the given (polynomial) spin Hamiltonian $\mathcal{H}$ on $\mathbb{C}^{2j+1}$, one has to replace $\mathcal{J}_+, \mathcal{J}_-$, and $\mathcal{J}_3$ by $\mathcal{J}_+/\sqrt{2j}, \mathcal{J}_-/\sqrt{2j}$, and $\mathcal{J}_3 + j\mathbb{1}$, respectively. If $\mathcal{H}_j = \int_{\mathbb{C}} d^2z\, h_j(z)|z\rangle\langle z|$ denotes the resulting operator, one then finds the relation

$$\lim_{j\to\infty} \frac{\pi}{2j} \langle z/\sqrt{2j}|\, e^{-it\mathcal{H}_j}\, |z'/\sqrt{2j}\rangle = \langle\!\langle z|\, e^{-it\mathsf{H}}\, |z'\rangle\!\rangle \tag{4}$$

where $|z\rangle\!\rangle \in \mathrm{L}^2(\mathbb{R})$ is a normalized canonical coherent vector and the Hamiltonian H on $\mathrm{L}^2(\mathbb{R})$, the Hilbert space of Lebesgue square-integrable complex-valued functions on the real line $\mathbb{R}$, is defined by $\mathsf{H} := \int_{\mathbb{C}} \frac{d^2z}{\pi}\, \mathsf{h}(z)|z\rangle\!\rangle\langle\!\langle z|$ with $\mathsf{h}(z) := \lim_{j\to\infty} h_j(z/\sqrt{2j})$. By using (3) for the pre-limit expression in (4), suitably rescaling the Brownian bridge, and interchanging the order of the limits $j \to \infty$ and $\nu \to \infty$, one arrives at the path-integral formula

$$\begin{aligned} \langle\!\langle z|\, e^{-it\mathsf{H}}\, |z'\rangle\!\rangle = \pi \lim_{\nu\to\infty} e^{2t\nu} \int & d\mu^{(\nu)}_{z,0;z',t}(b) \\ & \times \exp\left\{ \frac{1}{2}\int_0^t ds\, \left[\dot{b}(s)b^*(s) - \dot{b}^*(s)b(s)\right] - i\int_0^t ds\, \mathsf{h}(b(s)) \right\}\ , \end{aligned} \tag{5}$$

in agreement with Eq. (1.3) in Ref. 2, see also Ref. 11.

- With regard to some of the symbolic path-integral expressions for spin systems frequently encountered in the literature, see for example Refs. 4-7, it might be illuminating to recognize certain formal similarities between these expressions and the above result (3). While the kinematical and dynamical terms in the exponents of all the corresponding path integrands are essentially the same, only the above result is based on a genuine path measure, namely $d\mu^{(\nu)}_{z,0;z',t}(b)\exp\{4(j+1)\nu\int_0^t ds(1+|b(s)|^2)^{-2}\}$, but requires taking the limit $\nu \to \infty$. Here, the Wiener type of measure $d\mu^{(\nu)}_{z,0;z',t}(b)$ is often symbolically written as $\delta^2 b\, \delta(b(0)-z)\delta(b(t)-z')\exp\{-\frac{1}{4\nu}\int_0^t ds|\,\dot{b}(s)|^2\}$ or similarly.

3 Sketch of the proof

Adapting ideas of Ref. 2, the key to the proof of (3) is to identify the spin Hilbert space $\mathbb{C}^{2j+1}$ with the $(2j+1)$-dimensional ground-state eigenspace $E_0(\mathrm{L}^2(\mathbb{R}^2))$ of the positive Schrödinger operator

$$R := \left(i\frac{\partial}{\partial z_1} + \frac{2(j+1)z_2}{1+|z|^2}\right)^2 + \left(i\frac{\partial}{\partial z_2} - \frac{2(j+1)z_1}{1+|z|^2}\right)^2 - \frac{4(j+1)}{(1+|z|^2)^2} \tag{6}$$

acting on the Hilbert space $\mathrm{L}^2(\mathbb{R}^2)$. The claimed dimensionality of R's ground-state projector E_0 follows from the Aharonov-Casher theorem on zero-energy eigenstates.[12] By Taylor expanding about $t = 0$ one may now check that the coherent representation of the spin time-evolution operator may be written as

$$\langle z|e^{-it\mathcal{H}}|z'\rangle = E_0 e^{-itE_0HE_0}(z, z')\,. \tag{7}$$

Here the right-hand side denotes the continuous integral kernel (in other words position representation) of the operator $E_0 \exp\{-itE_0HE_0\}$, where the bounded multiplication operator H on $\mathrm{L}^2(\mathbb{R}^2)$ is defined by $(H\psi)(z) := h(z)\psi(z)$ for all $\psi \in \mathrm{L}^2(\mathbb{R}^2)$. It then remains to show the pointwise identity

$$E_0 e^{-itE_0HE_0}(z, z') = \lim_{\nu\to\infty} e^{-t(\nu R + iH)}(z, z')\,, \tag{8}$$

since the pre-limit expressions in (8) and (3) coincide by the Feynman-Kac-Itô formula.[8,13] Finally, to prove (8) one uses the strong convergence $\lim_{\nu\to\infty} \exp\{-t\nu R\} = E_0$ (for all $t > 0$) in the Duhamel-Dyson-Phillips perturbation expansion of $\exp\{-t(\nu R + iH)\}(z, z')$. For details see Ref. 3.

References

1. R. P. Feynman, *Space-time approach to non-relativistic quantum mechanics*, Rev. Mod. Phys. **20**, 367-387 (1948)
2. I. Daubechies, J. R. Klauder, *Quantum-mechanical path integrals with Wiener measure for all polynomial Hamiltonians II*, J. Math. Phys. **26**, 2239-2256 (1985)
3. B. Bodmann, H.Leschke, S. Warzel, *A rigorous path integral for quantum spin using flat-space Wiener regularization*, preprint 1998
4. H. Kuratsuji, *Path integrals in the $SU(2)$ coherent state representation and related topics*, in: A. Inomata, H. Kuratsuji, C. C. Gerry, *Path integrals and coherent states of $SU(2)$ and $SU(1,1)$*, World Scientific, Singapore 1992, pp. 139-218
5. K. Funahashi, T. Kashiwa, S. Sakoda, K. Fujii, *Coherent states, path integral, and semi-classical approximation*, J. Math. Phys. **36**, 3232-3253 (1995)
6. E. A. Kochetov, *SU(2) coherent-state path integral*, J. Math. Phys. **36**, 4667-4679 (1995)
7. E. Ercolessi, G. Morandi, F. Napoli, P. Pieri, *Path integrals for spinning particles, stationary phase and the Duistermaat-Heckman theorem*, J. Math. Phys. **37**, 535-553 (1996)
8. B. Simon, *Functional integration and quantum physics*, Academic, New York 1979
9. L. C. G. Rogers, D. Williams, *Diffusions, Markov processes, and martingales*, vol. 2: Itô calculus, Wiley, Chichester 1987
10. F. T. Arecchi, E. Courtens, R. Gilmore, H. Thomas, *Atomic coherent states in quantum optics*, Phys. Rev. A **6**, 2211-2237 (1972)
11. J. R. Klauder, *Coherent state path integrals at (nearly) 40*, quant-ph/9810043 and this volume
12. H. L. Cycon, R. G. Froese, W. Kirsch, B. Simon, *Schrödinger operators*, Springer, Berlin 1987
13. K. Broderix, D. Hundertmark, H. Leschke, *Continuity properties of Schrödinger semi-groups with magnetic fields*, math-ph/9808004, to appear in Rev. Math. Phys.